






Viewing the Universe at Very High Energies

VERITAS (Mt. Hopkins, AZ)



Rene A. Ong

MIT Colloquium

02 March 2009

Outline

Scientific Motivation

- A “New Astronomy”
- Physicist’s Viewpoint
 - *Astrophysical TeV accelerators*
 $1 \text{ TeV} = 10^{12} \text{ eV}$
 - *Origin of Cosmic Rays, understanding AGN ...*
 - *Probes of new physics, cosmology.*

Experimental Technique


The VERITAS Project

- Description, performance.
- Highlights of results from 18 months.
- Science Program, overlap with Fermi Telescope.


Future

A New Astronomy


- Before 1940's – Astronomy only used visible light.
- New wavebands (radio, IR, X-ray, γ -ray) change our picture of the universe
 - Different spatial scales
 - Different time scales
 - Different emission processes
 - ▶ New physics
- Other messengers (cosmic rays, neutrinos, grav. waves)



TeV γ -rays




New Windows & New Messengers




The TeV γ -ray Sky - 1998

3 sources




The TeV γ -ray Sky - 2003

12 sources



The TeV γ -ray Sky - 2008




- Explosion in number of sources, and a variety of source classes..
- Much more information: imaging, spectra, light curves ...

(Almost all) discoveries made by Atmospheric Cherenkov Telescopes

A Wide Variety of Sources ...

Supernova Remnants




Shocks
Fermi Mechanism

Pulsars/PWN




NS dynamo
Winds

HMXBs (microquasars)




Accretion-powered jets,
colliding winds, or ...?

Active Galactic Nuclei




Massive BH
Jets

Gamma-Ray Bursts



Massive star collapse
Int./ext. shocks

Dark accelerators...




???


... and accelerators

Key Physics Issues


SNR




AGN




GRBs




Origin of cosmic rays




Cosmological γ -ray horizon



Tests of Lorentz invariance




Cold dark matter (WIMP) searches



Origin of Cosmic Rays = SNRs ?

Why (VHE) gamma rays?


- Unlike cosmic rays, *not deflected* by interstellar magnetic fields.
- *Tracers* of parent particle populations – those particles accelerated by shocks.



SNR Image (RXJ 1713-3946)

Accelerated electrons
→ **VHE γ -rays**


Up-scattering of soft photons



Inverse Compton Scattering


Accelerated protons
→ **VHE γ -rays**

Target interaction, π^0 decay




Target material

There is evidence for SNR acceleration of CRs, but case is far from settled.



Spectral Energy Distribution

Active Galaxies



AGN cartoon

Active Galactic Nuclei


- High-luminosity extragalactic objects
 - Probe properties of the universe at large distances
- Variable!
- Jets powered by accretion on to supermassive BH

So far, AGN observed in VHE γ -rays are generally:

- **Blazars**
 - Jets aligned with line of sight
- Nearby: $z < 0.25$, EBL cutoff.
- Soft spectrum $\Gamma > 3.0$.


But not all are like this !

Extragalactic Background Light (EBL)



Diffuse extragalactic background light (how much light since recombination?)

- Complements direct measurement in Optical, IR: *difficult*.
- Absorption signature in 50-1000 GeV band for distant sources.




Red shifted
stellar light

Red shifted
dust light

Search for Cold Dark Matter



Hypothesis: DM = WIMPs

- Indirect detection of WIMP annihilation $\rightarrow \gamma, \nu$ etc.




Target regions with:

- Favorable DM distributions.
- Large mass/light ratio.



Complementary approach to direct detection & LHC.


Experimental Technique





Effective area
= light pool size
= 10^5 m^2 !!!

Whipple 10m γ -ray Telescope


- The Whipple 10m (1968 -).
- Pioneered use of Imaging.
T. Weekes et al.
- Made first source detections.




gamma ray?




cosmic ray?



Stereoscopy: Telescope Arrays




Major VHE Telescopes



Fermi

Multi-messenger Astronomy



VERITAS



Collaboration of ~80 scientists.
15 Institutions in U.S., Canada,
U.K., and Ireland.

Detector Design:

- Four 12m telescopes.
- 500 pixel cameras (3.5°).
- Site in southern Az (1300m).

Performance:

- Energy threshold ~ 100 GeV.
- Ang. resolution $\sim 4\text{-}6'$.
- Detect Crab Nebula in $\sim 45\text{s}$.

**Very Energy Radiation Imaging
Telescope Array System (VERITAS)**

Bumps in the Road



VERITAS: Mt. Hopkins, AZ



U.S.:

Adler Planetarium
Argonne National Lab
Barnard College
DePauw Univ.
Grinnell College
Iowa State Univ.
Purdue Univ.
Smithsonian

Univ. of California, Los Angeles
Univ. of California, Santa Cruz
Univ. of Chicago
Univ. of Delaware
Univ. of Iowa
Univ. of Massachusetts
Univ. of Utah
Washington Univ., St. Louis

Canada:

McGill Univ.

U.K.:

Leeds Univ.

Ireland:

Cork Inst. Tech.
Galway-Mayo Inst. Tech.
Nat. Univ. Ireland, Galway
Univ. College Dublin

+ ~25 Associate Members

Telescope Layout





T3
Fall 2006

T2
Spring 2006

T4
Spring 2007

T1
Jan 2005





7km

108m

T2
Spring 2006

85m

T3
Fall 2006

82m

T4
Spring 2007

T1
Jan 2005

35m

Relocating T1

T1
Dec 2009



95m



T2
Spring 2006

85m



T3
Fall 2006

108m


82m

80m

T4
Spring 2007

35m

Sensitivity Improvement ~ 20%




A VERITAS Telescope



12m reflector, f1.0 optics



350 Mirror Facets





500 pixel Camera

VERITAS Data Acquisition




- PMTs digitized with 500 MHz sampling FADCs
 - 24 samples/channel.
 - <6% deadtime @ 250 Hz.



Telescopes/cameras/electronics
meet all design specifications.

Four-Telescope Event




Crab Nebula – Now a Calibration !

VERITAS Sensitivity:


1 Crab 45s (5σ)

5% Crab ~2.5 hr

1% Crab ~40 hrs





Angular resolution 3'-6'
Pointing accuracy < 75''





Energy spectrum



VERITAS First-Year Results




XRB LSI +61




Discovery of SNR IC 443




SNR Cassiopeia A




M87 Radio Galaxy




Blazar 1ES1218+30
z=0.182, 2nd most distant VHE source



Discovery of Blazar W Comae
z = 0.102




Discovery of Blazar 1ES 0806
z = 0.138



Discovery of Blazar 3C 66A, z = 0.444
Most distant source ?

IC443: The Case For Hadronic Emission?

Excess Map (smoothed)



(1) Zhang, L. and Fang, J. Ap. J. **675** L21 (2008).
(2) Torres, D. F. et al. arXiv:0804.2526.




- VERITAS/MAGIC co-discovery, 2007.
- Overlap with CO indicating molecular cloud along line of sight.
- Maser emission suggests SNR shock interacting with cloud.
- TeV emission could be

➔ **CR-induced pion production in cloud** (1,2)


New Blazars with VERITAS

- ❑ W Comae: VERITAS discovery
 - ➔ **2 flares, 1 taken in moonlight.**
 - ➔ **First IBL discovered at VHE.**
 - ➔ **Simultaneous Swift data.**




Two AGN in the same field.

Astrophysical Journal, 684, L73, 2008



Light curve



Broad-band SED, EC preferred.



New Blazars with VERITAS II

3C66A

- Distant? $Z=0.444$ based on a single emission line (TeV spectrum raises questions).
- VERITAS excludes 3C66B (in 2008) at 4.3σ .

Controversy:
Blazar
or
Radio
galaxy?

MAGIC
claims
3C66B
(@85% CL)
in 2007



Very soft spectrum

$$\Gamma = 4.1 \pm 0.4 \pm 0.6$$

If $z=0.444$, $\Gamma_{int} = 1.1$


Via Franceschini et al. 2008

Acciari et al., ApJ, in press, 2009; astro-ph/0901.4527.

Also, significant MWL paper with Fermi, Swift, XMM, optical, radio.

Dark Matter Search: Dwarf Galaxies

Recent VERITAS observations:
 Draco, Ursa Minor, **Willman I**



Whipple 10m
 Ursa Minor constraint

VERITAS (and 15 hr data)
Willman I constraints
 derived from
 Strigari et al. 2007

$\langle\sigma v\rangle$:
 Thermal average of product
 σ : WIMP self-annihilation x-sec
 v : WIMP velocity


Minimal Supersymmetric extensions to Standard Model (MSSM) allowed by WMAP





What's next for VERITAS ?

LOTS!

- New Results: to be announced this summer.
- Observing: we are in 2nd year of 5+ year program.
- Fermi Gama-Ray Space Telescope overlap.
- Spectra and modelling: source mechanisms.
- MWL studies: radio, optical, X-ray, γ -ray.
- Upgrade possibilities: e.g. new cameras, triggers.
- ...

VERITAS Sky Survey



-  SNR/PWN
-  TeV Unidentified
-  EGRET (GeV)
-  X-ray binaries


Other possible source types:
star clusters/star-forming regions, Wolf-Rayet stars

..or the completely unexpected !


- 2 year project, covering 150 deg².
- Ambitious---originally intended a larger region, but getting enough time is challenging (region best visible in July/August moonsoon season).
- 150 hrs data taken so far, results this summer.

Fermi GST

Large Area Telescope (LAT)



LAT images the sky one photon at a time:
 γ -ray converts in LAT to an electron and a positron ; direction and energy of these particles tell us the direction and energy of the photon



Fermi-LAT science objectives

> 2000 AGNs

blazars and radiogal = $f(\theta, z)$
evolution $z < 5$
Sag A*

10-50 GRB/year

GeV afterglow
spectra to high energy

γ -ray binaries

Pulsar winds
 μ -quasar jets

Cosmic rays and clouds

acceleration in Supernova remnants
OB associations
propagation (Milky Way, M31, LMC, SMC)
Interstellar mass tracers in galaxies



Possibilities

starburst galaxies
galaxy clusters
measure EBL
unIDs

Dark Matter

neutralino lines
sub-halo clumps

Pulsars


emission from radio and X-ray pulsars
blind searches for new Gemingas
magnetospheric physics
pulsar wind nebulae





Launch from Cape Canaveral Air Station
11 June 2008 at 12:05PM EDT.




First Light FGST-LAT


- ~4-day First Light exposure, June 30 – July 3, 2008.
- Orthographic projection.
- Comparable to EGRET on CGRO!







Fermi γ -ray Sky (Feb 2009)







VERITAS Source Catalog (18 mo)



<p>Outside RSS iWUFS</p> <p>Other MacOS: Dashboard Widget</p>	<p style="text-align: center;">The Astronomer's Telegram for reporting and commenting on new astronomical observations</p> <p style="text-align: center;">Post a New Telegram Submit Information Minor Software Telegram Index</p> <p style="text-align: center;">Register To Post Email and RSS Subscriptions Forgot your password?</p> <p style="text-align: center;">Present Time: 1 Mar 2009; 7:00 UT</p> <p style="text-align: center;">[Previous Next]</p> <p>VERITAS Discovery of VHE Gamma-Ray Emission from BL Lac object RGB J0710+591</p> <p>ATel #1941: Read One (UCI-A) for the VERITAS Collaboration on 25 Feb 2009; 0:32 UT</p> <p><i>Distributed as an Instant Email Notice (Request for Observations)</i> Password Certification: Philip Kazarek (philip-kazarek@uiowa.edu)</p> <p>Subjects: >GeV, Request for Observations, AGN</p> <p>The VERITAS collaboration reports the discovery of very high energy (VHE; E>100 GeV) gamma-ray emission from the BL Lac object RGB J0710+591 (z=0.125). This new VHE source was observed for ~18 hours good-quality live time between 3 January and 23 February 2009 (UT) with the VERITAS atmospheric-Cherenkov telescope array. Preliminary analysis of these data yields a detection of ~100 gamma-rays from RGB J0710+591 corresponding to a significance of >5 standard deviations. The VHE flux is ~1.6% of the Crab above 300 GeV, and there is no evidence of flux variability. VERITAS will continue to observe RGB J0710+591 and contemporaneous multi-wavelength observations of this blazar are encouraged.</p>
---	---

FUTURE

Next 5-10 years will be exciting period for this field:

VERITAS will survey the northern TeV sky with great sensitivity, complementing:

Fermi-LAT (GeV, in space)

HESS (TeV, S. Hemisphere)


IceCube (ν , South Pole)

Auger S (UHECR, S. Hemisphere)

Farther in the future:

- Astrophysics at GeV & TeV energies with large km² Cherenkov Telescope arrays.

The Next Generation



- Populations of fainter sources we have yet to probe.
- TeV source confusion may be starting to be an issue.
 - **very deep observations to get morphology, disentangle sources**
- Need larger source populations to get away from source idiosyncrasies.
- Next few years with Fermi may help to answer questions
 - **...but not completely.**
 - **...and we will have many new sources !**


AGIS (Advanced Gamma Imaging System)

Large (1 km²) array.

- ~50-75 telescopes, aperture 8-20m.
- \$100-150M class observatory.


Much more sensitive than FGST/VERITAS..
 APS White Paper study, collaboration formed.

CTA (Europe) – considerable momentum.



Transition to Fermi Regime (GRBs, etc.)

Spectral cutoffs (acceleration mechanisms)



8-10m telescope
 Wide-field



“Ground-Based Gamma Ray Astronomy, Towards the future”, Oct. 20-21 2005, Malibu, CA (UCLA)

“Ground-Based Gamma Ray Astronomy, Towards the future”, May 11-12 2006, Santa Fe, NM (LANL)

**“Future of Very High Energy Gamma Ray Astronomy, Towards the future”, May 13-14 2007, Chicago, IL, (UofChicago) –
Formation of AGIS R&D group**

**Collaboration meeting, June 27-28 2008, UCLA,
Formation of AGIS collaboration**

Institutions:

**ADLER
ANL
Barnard
Delaware
IAFE
Iowa State
LANL
McGill
Penn State
Purdue**

**SAO
Stanford/KIPAC
UNAM
UC, Los Angeles
UC, Santa Cruz
U. Chicago
U. Iowa
Utah
Yale
Washington U.**

**Wide-Field
Schwarzschild-Coudet Telescope**



Summary

- HE γ -rays provide unique tests of the limits of physical laws. Probe astrophysics in regimes not yet explored. Possibility for discovery of physics beyond our standard models.
- Exciting discoveries of many, unexpected sources of VHE gamma-rays. But still, most of the sky remains unexplored.
 - **VERITAS and Fermi are now both operational and getting exciting results.**
- New Astronomy of TeV γ -rays (and neutrinos, grav. waves) should reveal many surprises over the next 10 years.

“The real voyage of discovery consists, not in seeking new landscapes, but in having new eyes.”

Marcel Proust (1871-1922)

